arXiv:1603.02065v1 [math.CA] 22 Feb 2016

A CLASS OF FUNCTIONAL EQUATIONS ON MONOIDS

BOUIKHALENE BELAID AND ELQORACHI ELHOUCIEN

Abstract. In [8] B. Ebanks and H. Stetkær obtained the solutions of the functional equation $f(xy) - f(\sigma(y)x) = g(x)h(y)$ where σ is an involutive automorphism and f, g, h are complex-valued functions, in the setting of a group G and a monoid M. Our main goal is to determine the complex-valued solutions of the following more general version of this equation, viz $f(xy) - \mu(y)f(\sigma(y)x) = g(x)h(y)$ where $\mu : G \longrightarrow \mathbb{C}$ is a multiplicative function such that $\mu(x\sigma(x)) = 1$ for all $x \in G$. As an application we find the complex-valued solutions (f, g, h) on groups of the equation $f(xy) + \mu(y)g(\sigma(y)x) = h(x)h(y)$.

1. Introduction

We recall that a semigroup S is a set equipped with an associative operation. We write the operation multiplicatively. A monoid is a semigroup M with identity element that we denote e. A function $\chi : S \longrightarrow \mathbb{C}$ is said to be multiplicative if $\chi(xy) = \chi(x)\chi(y)$ for all $x, y \in S$.
Let S be a semigroup and $\sigma : G \longrightarrow G$ an involutive homomorphism, that is $\sigma(xy) = \sigma(x)\sigma(y)$ and $\sigma(\sigma(x)) = x$ for all $x, y \in G$.
The complex-valued solutions of the following variant of d'Alembert's functional equation

$$f(xy) + f(\sigma(y)x) = 2f(x)f(y),\ x, y \in S. \tag{1.1}$$

was determined by Stetkær [14]. They are the functions of the form

$$f(x) = \frac{\chi + \chi \circ \sigma}{2},$$

where $\chi : G \longrightarrow \mathbb{C}$ is multiplictive.
In the same year, Ebanks and Stetkær [8] obtained on monoids the complex-valued solutions of the following d'Alembert's other functional equation

$$f(xy) - f(\sigma(y)x) = g(x)h(y),\ x, y \in S. \tag{1.2}$$

This functional equation contains, among others, an equation of d'Alembert [2, 3, 4]

$$f(x + y) - f(x - y) = g(x)h(y),\ x, y \in \mathbb{R} \tag{1.3}$$

the solutions of which are known on abelian groups, and a functional equation

$$f(x + y) - f(x + \sigma(y)) = g(x)h(y),\ x, y \in G \tag{1.4}$$

studied by Stetkær [12, Corollary III.5] on abelian groups G.
There are various ways of extending functional equations from abelian groups to non-abelian groups. The μ-d'Alembert functional equation

$$f(xy) + \mu(y)f(x\tau(y)) = 2f(x)f(y),\ x, y \in G \tag{1.5}$$

which is an extension of d'Alembert functional equation

$$f(xy) + f(x\tau(y)) = 2f(x)f(y),\ x, y \in G, \tag{1.6}$$

where τ is an involutive of G, is closely related to pre-d'Alembert functions. It occurs in the literature. See Parnami, Singh and Vasudeva [10], Davison [5, Proposition 2.11], Ebanks and Stetkær [6], Stetkær [15, Lemma IV.4] and Yang [16, Proposition 4.2]. The functional equation (1.5) has been treated systematically by Stetkær [15]. The non-zero solutions of (1.5) are the normalized traces of certain representations of G on $\mathbb{C}^2$. Davison proved this via his work [5] on the pre-d'Alembert functional equation on monoids.
The variant of Wilson's functional equation

$$f(xy) + \mu(y)f(\tau(y)x) = 2f(x)g(y),\ x, y \in G; \tag{1.7}$$

$$f(xy) + \mu(y)f(\sigma(y)x) = 2f(x)g(y),\ x, y \in G \tag{1.8}$$

with $\mu \neq 1$ was recently studied on groups by Elqorachi and Redouani [9].
The complex-valued solutions of equation (1.7) with $\tau(x) = x^{-1}$ and $\mu(x) = 1$ for all $x \in G$ are obtained on groups by Ebanks and Stetkær [6].
The present paper complements and contains the existing results about (1.2) by finding the solutions f, g, h of the extension

$$f(xy) - \mu(y)f(\sigma(y)x) = g(x)h(y),\ x, y \in S \tag{1.9}$$

of it to monoids that need not be abelian, because on non-abelian monoids the order of factors matters and involutions and involutive automorphisms differ.
As in [8] one of the main ideas is to relate the functional equation (1.2) to a sine substraction law on monoids. In our case we need the solutions of the following version of the sine subtraction law

$$\mu(y)f(x\sigma(y)) = f(x)g(y) - f(y)g(x),\ x, y \in S. \tag{1.10}$$

These results are obtained in Theorem 2.1. We need also the solutions of equation (1.8) on monoids. There are not in the literature, but we derived them in Theorem 3.2. In section 4 we obtain on of the main results about (1.9). Furthermore, as an application we find the complex-valued solutions (f, g, h) of the functional equation

$$f(xy) + \mu(y)g(\sigma(y)x) = h(x)h(y),\ x, y \in G \tag{1.11}$$

on groups and monoids in terms of multiplicative and additive functions.

1.1. **Notation and preliminary matters.** Throughout this paper G denotes a group and S a semigroup. A monoid is a semigroup M with an identity element that we denote e. The map $\sigma : S \longrightarrow S$ denotes an involutive automorphism. That it is involutive means that $\sigma(\sigma(x)) = x$ for all $x \in S$. The mapping $\mu : S \longrightarrow \mathbb{C}$ is a multiplicative function such that $\mu(x\sigma(x)) = 1$ for all $x \in S$. If $\chi : S \longrightarrow \mathbb{C}$ is a multiplicative function such that $\chi \neq 0$, then $I_\chi = \{x \in S \mid \chi(x) = 0\}$ is either empty or a proper subset of S. Furthermore, I_χ is a two-sided ideal in S if not empty and $S \backslash I_\chi$ is a subsemigroup of S. If S is a topological space, then we let $\mathcal{C}(S)$ denote the algebra of continuous functions from S into $\mathbb{C}$.
For later use we need the following results. The next proposition corresponds to Lemma 3.4 in [8].

Proposition 1.1. *Let S be a semigroup, and suppose $f, g : S \longrightarrow \mathbb{C}$ satisfy the sine addition law*

$$f(xy) = f(x)g(y) + f(y)g(x),\ x, y \in S \tag{1.12}$$

with $f \neq 0$. Then there exist multiplicative functions $\chi_1, \chi_2 : G \longrightarrow \mathbb{C}$ such that

$$g = \frac{\chi_1 + \chi_2}{2}.$$

Additionally we have the following
i) If $\chi_1 \neq \chi_2$, then $f = c(\chi_1 - \chi_2)$ for some constant $c \in \mathbb{C} \setminus \{0\}$.
ii) If $\chi_1 = \chi_2$, then letting $\chi := \chi_1$ we have $g = \chi$. If S is a semigroup such that $S = \{xy \in : x, y \in S\}$ (for instance a monoid), then $\chi \neq 0$.
If S is a group, then there is an additive function $A : S \longrightarrow \mathbb{C}$, $A \neq 0$, such that $f = \chi A$.
If S is a semigroup which is generated by its squares, then there exists an additive function $A : S \setminus I_\chi \longrightarrow \mathbb{C}$ for which

$$f(x) = \begin{cases} \chi(x)A(x) & for\ x \in S \setminus I_\chi \\ 0 & for\ x \in I_\chi. \end{cases}$$

Furthermore, if S is a topological group, or if S is a topological semigroup generated by its squares, and $f, g \in \mathcal{C}(S)$, then $\chi_1, \chi_2, \chi \in \mathcal{C}(S)$. In the group case $A \in \mathcal{C}(S)$ and in the second case $A \in \mathcal{C}(S \setminus I_\chi)$.

2. μ-sine subtraction law on a group and on a monoid

In this section we deal with a new version of the sine subtraction law

$$\mu(y)k(x\sigma(y)) = k(x)l(y) - k(y)l(x),\ x, y \in S, \tag{2.1}$$

where k, l are complex valued functions and μ is a multiplicative function. The new feature is the introduction of the function μ. We shall say that k satisfies the μ-sine subtraction law with companion function l. If S is a topological semigroup and k, l satisfy (2.1) such that $k \neq 0$ and k is a continuous function then l is also a continuous function. In the case where $\mu = 1$ and G is a topological group, the functional equation (2.1) was solved in [8].
Here we focus exclusively on (2.1), and we include nothing about other extensions of the cosine, sine addition and subtraction laws. The next theorem is the analogue of Theorem 3.2 in [8].

Theorem 2.1. *Let G be a group and let $\sigma : G \longrightarrow \mathbb{C}$ be a involutive automorphism. Let $\mu : G \longrightarrow \mathbb{C}$ be a multiplicative function such that $\mu(x\sigma(x)) = 1$ for all $x \in G$. The solution $k, l : G \longrightarrow \mathbb{C}$ of the μ-sine subtraction law (2.1) with $k \neq 0$ are the following pairs of functions, where $\chi : G \longrightarrow \mathbb{C} \setminus \{0\}$ denotes a character and $c_1 \in \mathbb{C}$, $c_2 \in \mathbb{C} \setminus \{0\}$ are constants.*
i) If $\chi \neq \mu\, \chi \circ \sigma$, then

$$k = c_2 \frac{\chi - \mu\, \chi \circ \sigma}{2},\ l = \frac{\chi + \mu\, \chi \circ \sigma}{2} + c_1 \frac{\chi - \mu\, \chi \circ \sigma}{2}.$$

ii) If $\chi = \mu\, \chi \circ \sigma$, then

$$k = \chi A,\ \ l = \chi(1 + c_1 A)$$

where $A : G \longrightarrow \mathbb{C}$ is an additive function such that $A \circ \sigma = -A \neq 0$.
Furthermore, if G is a topological group and $k \in \mathcal{C}(G)$, then $l, \chi, \mu\, \chi \circ \sigma, A \in \mathcal{C}(G)$.

Proof. By interchanging x and y in (2.1) we get that $\mu(x)k(y\sigma(x)) = -\mu(y)k(x\sigma(y))$ for all $x, y \in G$. By setting $y = e$ we get that $k(x) = -\mu(x)k(\sigma(x))$ for all $x \in G$.

Using this, equation (2.1) and the fact that $\mu(x\sigma(x)) = 1$ we get for all $x, y \in G$ that

$$\begin{aligned}
& k(x)[l(y) - \mu(y)l(\sigma(y))] - k(y)[l(x) - \mu(x)l(\sigma(x))] \\
& = k(x)l(y) - \mu(y)k(x)l(\sigma(y)) - k(y)l(x) + \mu(x)k(y)l(\sigma(y)) \\
& = \mu(y)k(x\sigma(y) + \mu(xy)k(\sigma(x))l(\sigma(y)) - \mu(xy)k(\sigma(y))l(\sigma(x)) \\
= \quad & \mu(y)k(x\sigma(y)) + \mu(xy)[k(\sigma(x))l(\sigma(y)) - k(\sigma(y))l(\sigma(x))] \\
& = \mu(y)k(x\sigma(y)) + \mu(xy)\mu(\sigma(y))k(\sigma(x)y) \\
& = \mu(y)k(x\sigma(y)) + \mu(x)k(\sigma(x)y) \\
& = mu(y)k(x\sigma(y)) - \mu(x)\mu(\sigma(x)y)k(x\sigma(y)) \\
& = \mu(y)k(x\sigma(y)) - \mu(y)k(x\sigma(y)) \\
& = 0.
\end{aligned}$$

So that we get for all $x, y \in G$ that

$$k(x)[l(y) - \mu(y)l(\sigma(y))] = k(y)[l(x) - \mu(x)l(\sigma(x))]. \tag{2.2}$$

By using some ideas from [8] we let $l^+(x) = \frac{l(x)+\mu(x)l(\sigma(x))}{2}$ and $l^-(x) = \frac{l(x)-\mu(x)l(\sigma(x))}{2}$ for all $x \in G$. We have $l = l^+ + l^-$, $l^+(\sigma(x)) = \mu(\sigma(x))l^+(x)$ and $l^-(\sigma(x)) = -\mu(\sigma(x))l^-(x)$ for all $x \in G$. From (2.2) we have for all $x, y \in G$ that $k(x)l^-(y) = k(y)l^-(x)$. Since $k \neq 0$ there exists an $x_0 \in G$ such that $k(x_0) \neq 0$. Thus we have $l^-(y) = \frac{l^-(x_0)}{k(x_0)}k(y) = ck(y)$ for all $y \in G$. Then $l^- = ck$. On the other hand by replacing l by $l^+ + l^-$ in (2.1) and by using $k(x)l^-(y) = k(y)l^-(x)$ we get that

$$\mu(y)k(x\sigma(y)) = k(x)l^+(y) - k(y)l^+(x),\ x, y \in G \tag{2.3}$$

Replacing y by $\sigma(y)$ in (2.3) we get that

$$k(xy) = k(x)l^+(y) + k(y)l^+(x),\ x, y \in G. \tag{2.4}$$

According to Proposition 1.1 we have
either i) or ii) $k = c_1(\chi_1 - \chi_2)$ and $l^+ = \frac{\chi_1+\chi_2}{2}$ where $c_1 \in \mathbb{C} \setminus \{0\}$. Since $k(x) = -\mu(x)k(\sigma(x))$ and $l^+(x) = \mu(x)l^+(\sigma(x))$ for all $x \in G$ we get that $c_1(\chi_1 - \chi_2) = -c_1\mu(\chi_1 \circ \sigma - \chi_2 \circ \sigma)$ and $\mu(\chi_1 \circ \sigma + \chi_2 \circ \sigma) = \chi_1 + \chi_2$. Then $\chi_2 = \mu\chi_1 \circ \sigma$, $l^+ = \frac{\chi_1+\mu\chi_1\circ\sigma}{2}$ and $k = c_1\frac{\chi_1-\mu\chi_1\circ\sigma}{2}$. Since $l^- = ck = cc_1\frac{\chi_1-\mu\chi_1\circ\sigma}{2} = c_2\frac{\chi_1-\mu\chi_1\circ\sigma}{2}$ where $c_2 \in \mathbb{C} \setminus \{0\}$. By using the fact $l = l^- + l^+$ we get (i).
ii) we have $l^+ = \chi$ and $k = \chi A$. Since $l^+(\sigma(x)) = \mu(\sigma(x))l^+(x)$ for all $x \in G$ we get that $\chi = \mu\chi \circ \sigma$. From $k(x) = -\mu(x)k(\sigma(x))$ for all $x \in G$ we get that $A = -A \circ \sigma$. By using the fact $l = l^- + l^+ = c\chi A + \chi = \chi(cA + 1)$. This completes the proof. □

In the next proposition we extend Theorem 2.1 to monoids.

Proposition 2.2. *Let M be a monoid and let $\sigma : M \longrightarrow M$ be a involutive automorphism. Let $\mu : M \longrightarrow \mathbb{C}$ be a multiplicative function such that $\mu(x\sigma(x)) = 1$ for all $x \in M$. The solution $k, l : M \longrightarrow \mathbb{C}$ of the μ-sine subtraction law (2.1) with $k \neq 0$ are the following pairs of functions, where $\chi : M \longrightarrow \mathbb{C} \setminus \{0\}$ denotes a multiplicative function and $\chi(e) = 1$ and $c_1 \in \mathbb{C}$, $c_2 \in \mathbb{C} \setminus \{0\}$ are constants :*
i) If $\chi \neq \mu\,\chi \circ \sigma$, then

$$k = c_1\frac{\chi - \mu\,\chi \circ \sigma}{2},\ l = \frac{\chi + \mu\,\chi \circ \sigma}{2} + c_2\frac{\chi - \mu\,\chi \circ \sigma}{2}.$$

ii) If $\chi = \mu\,\chi\circ\sigma$ and S is generated by its squares, then

$$k(x) = \chi(x)A(x),\quad l(x) = \chi(x)(1 + c_1A(x))\ \forall x \in M\backslash I_\chi$$
$$l(x) = k(x) = 0,\quad \forall x \in I_\chi$$

where $A : M \longrightarrow \mathbb{C}\setminus\{0\}$ is an additive function such that $A\circ\sigma = -A \neq 0$. Furthermore, if M is a topological monoid, and $k \in \mathcal{C}(M)$, then $l, \chi, \mu\chi\circ\sigma \in \mathcal{C}(M)$ and $A \in \mathcal{C}(M\backslash I_\chi)$.

Proof. By the same way in Proposition 3.6 in [8] and by using Theorem 2.1 and the Proposition 1.1 we get the proof. □

3. A variant of Wilson's functional equation on monoids

The solutions of the functional equation (1.8) on groups are obtained in [9]. More precisely, we have the following theorem.

Theorem 3.1. *Let G be a group, let $\sigma : G \longrightarrow G$ a homomorphism such that $\sigma\circ\sigma = I$, where I denotes the identity map, and $\mu : G \longrightarrow \mathbb{C}$ be a multiplicative function such that $\mu(x\sigma(x)) = 1$ for all $x \in G$. The solutions f, g of the functional equation (1.8) are the following pairs of functions, where $\chi : G \longrightarrow \mathbb{C}$ denotes a multiplicative function and $c, \alpha \in \mathbb{C}^*$*
(i) $f = 0$ and g arbitrary.
(ii) $g = \frac{\chi+\mu\chi\circ\sigma}{2}$ and $f = \alpha g$.
(iii) $g = \frac{\chi+\mu\chi\circ\sigma}{2}$ and $f = (c+\alpha/2)\chi - (c-\alpha/2)\chi\circ\sigma$ with $(\mu-1)\chi = (\mu-1)\chi\circ\sigma$.
(iv) $g = \chi$ and $f = \chi(a+\alpha)$, where where $\chi = \mu\,\chi\circ\sigma$ and a is an additive map which satisfies $a\circ\sigma + a = 0$.

We shall now extend this result to monoids.

Theorem 3.2. *Let M be a monoid, let $\sigma : M \longrightarrow M$ a homomorphism such that $\sigma\circ\sigma = I$, where I denotes the identity map, and $\mu : M \longrightarrow \mathbb{C}$ be a multiplicative function such that $\mu(x\sigma(x)) = 1$ for all $x \in M$. The solutions f, g of the functional equation (1.8) are the following pairs of functions, where $\chi : M \longrightarrow \mathbb{C}$ denotes a multiplicative function and $c, \alpha \in \mathbb{C}^*$*
(i) $f = 0$ and g arbitrary.
(ii) $g = \frac{\chi+\mu\chi\circ\sigma}{2}$ and $f = \alpha g$.
(iii) $g = \frac{\chi+\mu\chi\circ\sigma}{2}$ and $f = (c+\alpha/2)\chi - (c-\alpha/2)\chi\circ\sigma$ with $(\mu-1)\chi = (\mu-1)\chi\circ\sigma$.
(iv) $g = \chi$, $f_e(xy) = f_e(x)\chi(y) + f_e(y)\chi(x)$ for all $x, y \in S$ and with $f = f_e + f(e)\chi$. Furthermore, if M is a monoid which is generated by its squares, then $\chi = \mu\,\chi\circ\sigma$, there exists an additive function $a : M\backslash I_\chi \longrightarrow \mathbb{C}$ for which $a\circ\sigma + a = 0$,

$$f(x) = \begin{cases} \chi(x)(a(x)+\alpha) & for\ x \in M\setminus I_\chi \\ 0 & for\ x \in I_\chi \end{cases}$$

Indeed, If M is a topological group, or M is a topological monoid generated by its squares, $f, g, \mu \in C(M)$, and $\sigma : M \longrightarrow M$ is continuous, then $\chi \in C(M)$. In the group case $a \in C(M)$ and in the second case $a \in C(M\backslash I_\chi)$.

Proof. Verifying that the stated pairs of functions constitute solutions consists of simple computations. To see the converse, i.e., that any solution f, g of (1.8) is contained in one of the cases below, we will use [8, Lemma 3.4] and [9, Theorem 3.1]. All, except the last paragraphs of part (iv) and the continuity statements, is in Theorem 3.1 in [9]. Now, we assume that M is a monoid generated by its

squares. We use the notation used in the proof of Theorem 3.1 in [9], in particular for the last paragraphs of part (iv) we have

$$f_e(xy) = f_e(x)\chi(y) + f_e(y)\chi(x) \tag{3.1}$$

for all $x, y \in S$ and with $f = f_e + f(e)\chi$. So, from [9, Lemma 3.4] we get $f(x) = 0 + f(e)\chi(x) = 0 + f(e)0 = 0$ if $x \in I_\chi$ and $f(x) = \chi(x)(a(x) + f(e))$ if $x \in S\backslash I_\chi$ and where a is an additive function of $S\backslash I_\chi$.
Now, we will verify that $\chi = \mu\ \chi \circ \sigma$ and $a \circ \sigma = -a$. Since f, g are solution of equation (1.8) we have

$$f(xy) + \mu(y)f(\sigma(y)x) = 2f(x)\chi(y) \tag{3.2}$$

for all $x, y \in M$. By using the new expression of f and the fact that I_χ is an ideal, we get after an elementary computation that $f(xy) = f(yx)$ for all $x, y \in M$. So, equation (3.2) can be written as follows

$$f(xy) + \mu(y)f(x\sigma(y)) = 2f(x)\chi(y),\ x, y \in M. \tag{3.3}$$

By replacing y by $\sigma(y)$ in (3.3) and multiplying the result obtained by $\mu(y)$ we get

$$f(xy) + \mu(y)f(x\sigma(y)) = 2f(x)\mu(y)\chi(\sigma(y)),\ x, y \in M. \tag{3.4}$$

Finally, by comparing (3.3), (3.4) and using $f \neq 0$ we get $\chi(y) = \mu(y)\chi(\sigma(y))$ for all $y \in M$.
By Substituting the expression of f into (3.3) and using $\chi(y) = \mu(y)\chi(\sigma(y))$ and $\mu(y\sigma(y)) = 1$ for all $y \in M$, we find after reduction that $\chi(x)\chi(y)[a(y)+a(\sigma(y))] = 0$ for all $x, y \in M\backslash I_\chi$. Since $\chi \neq 0$ we get $a(y) + a(\sigma(y)) = 0$ for all $y \in M\backslash I_\chi$.
For the topological statement we use [13, Theorem 3.18(d)]. This completes the proof. □

4. Solutions of (1.9) on groups and monoids

In this section we solve the functional equation (1.9) on monoids. In the next proposition we show that if (f, g, h) is a complex-valued solution of equation (1.9), then h satisfies the μ-sine subtraction law.

Proposition 4.1. *Let M be a monoid, let σ be a involutive automorphism on S, let μ be a multiplicative function on M such that $\mu(x\sigma(x)) = 1$ for all $x \in M$. Suppose that $f, g, h : M \longrightarrow \mathbb{C}$ satisfy the functional equation (1.9). Suppose also that $g \neq 0$ and $h \neq 0$. Then*
i) $h(\sigma(x)) = -\mu(\sigma(x))h(x))$ for all $x \in M$.
ii) $h(xy) = h(yx)$ for all $x, y \in M$.
3i) h satisfies the μ-sine subtraction law (2.1).
4i) If $g(e) = 0$, then $g = bh$ for some $b \in \mathbb{C} \setminus \{0\}$.
5i) If $g(e) \neq 0$, then h satisfies the μ-sine subtraction law with companion function $\frac{g}{g(e)}$.
Moreover, if M is a topological monoid, and $h \neq 0$ is continuous, then the companion function is also continuous.

Proof. We follow the path of the proof of Proposition 3.1 in [8].
By substituent $(x, yz), (\sigma(y), \sigma(z)x)$ and $(z, \sigma(xy))$ and $(z, \sigma(xy))$ in (1.9) we obtain

$$f(xyz) - \mu(yz)f(\sigma(yz)x) = g(x)h(yz), \tag{4.1}$$

$$f(\sigma(yz)x) - \mu(\sigma(z)x)f(z\sigma(xy)) = g(\sigma(y))h(\sigma(z)x), \tag{4.2}$$

$$f(z\sigma(xy)) - \mu(\sigma(xy))f(xyz) = g(z)h(\sigma(xy)). \tag{4.3}$$

By multiplying (4.1) by $\mu(\sigma(xy))$ we obtain that

$$\mu(\sigma(xy))f(xyz) - \mu(\sigma(x)z)f(\sigma(yz)x) = \mu(\sigma(xy))g(x)h(yz). \tag{4.4}$$

By adding (4.3) and (4.4) we obtain

$$f(z\sigma(xy)) - \mu(\sigma(x)z)f(\sigma(yz)x) = g(z)h(\sigma(xy)) + \mu(\sigma(xy))g(x)h(yz). \tag{4.5}$$

By multiplying (4.5) by $\mu(\sigma(z)x)$ we obtain

$$\mu(\sigma(z)x)f(z\sigma(xy)) - f(\sigma(yz)x) = \mu(\sigma(z)x)g(z)h(\sigma(xy)) + \mu(\sigma(zy))g(x)h(yz). \tag{4.6}$$

By adding (4.6) and (4.2) we obtain

$$0 = g(\sigma(y))h(\sigma(z)x) + \mu(\sigma(z)x)g(z)h(\sigma(xy)) + \mu(\sigma(zy))g(x)h(yz). \tag{4.7}$$

Setting x_0 such that $g(x_0) \neq 0$ and the fact that $\mu(x\sigma(x)) = \mu(x)\mu(\sigma(x)) = 1$ for all $x \in M$, we get that

$$h(yz) = \mu(y)g(\sigma(y))l(z) + g(z)l_1(y),\ y, z \in M \tag{4.8}$$

where l, l_1 are complex valued functions on M. Using (4.8) in (4.7) we obtain for all $x, y, z \in M$

$$g(x)g(\sigma(y))\{l_1(\sigma(z)) + \mu(\sigma(z))l(z)\} + g(x)g(z)\{\mu(\sigma(z))l(\sigma(y)) + \mu(\sigma(zy))l_1(y)\} + g(z)g(\sigma(y))\{\mu(\sigma(z))l(x) + \mu(\sigma(z)x)l_1(\sigma(x))\} = 0. \tag{4.9}$$

Putting $x = x_0$, $y = \sigma(x_0)$, the equation (4.9) becomes

$$l_1(\sigma(z)) + \mu(\sigma(z))l(z) = c\mu(\sigma(z))g(z),\ z \in M, \tag{4.10}$$

where $c \in \mathbb{C}$ is a constant. By putting (4.10) in (4.9) we obtain $3\mu(\sigma(z))cg(x)g(\sigma(y))g(z) = 0$ for all $x, y, z \in S$. Since $g \neq 0$ and $\mu(\sigma(z)) \neq 0$ it follows that $c = 0$ and then $l_1(\sigma(z)) = -\mu(\sigma(z))l(z)$ for all $z \in S$. So that equation (4.8) becomes

$$h(yz) = \mu(y)g(\sigma(y))l(z) - \mu(y)g(z)l(\sigma(y)),\ x, y, z \in S \tag{4.11}$$

Replacing (y, z) by $(\sigma(z), \sigma(y))$ in (4.11) we obtain

$$h(\sigma(zy)) = \mu(\sigma(zy))(\mu(y)g(z)l(\sigma(y)) - \mu(y)g(\sigma(y))l(z)) = -\mu(\sigma(zy))h(yz). \tag{4.12}$$

From which we obtain by putting $y = e$ that

$$h(\sigma(z)) = -\mu(\sigma(z))h(z),\ z \in M. \tag{4.13}$$

From (4.12) and (4.13) we get that h a central function i.e. $h(yz) = h(zy)$ for all $y, z \in M$.

Next we consider two cases :

First case : Suppose $g(e) = 0$. Let $z = e$ and $x = x_0$ in (4.7) give that

$$h(y) = -c\mu(y)g(\sigma(y)),\ y \in M \tag{4.14}$$

for some $c \in \mathbb{C} \setminus \{0\}$. By replacing y by $\sigma(y)$ in (4.14) we obtain that $h(\sigma(y)) = -c\mu(\sigma(y))g(y)$ for all $y \in M$. By using (4.14) we get that $g = \frac{1}{c}h = bh$ where $b = \frac{1}{c}$ and that

$$g(\sigma(z)) = -\mu(\sigma(z))g(z),\ z \in S. \tag{4.15}$$

Using (4.15) in (4.11) and setting $m = -bl$ we get

$$\begin{aligned} h(yz) &= \mu(y)g(\sigma(y))l(z) - \mu(y)g(z)l(\sigma(y)) \\ &= -\mu(y)\mu(\sigma(y))g(y)l(z) - \mu(y)g(z)l(\sigma(y)) \\ &= -g(y)l(z) - \mu(y)g(z)l(\sigma(y)) \\ &= -bh(y)l(z) - \mu(y)bh(z)l(\sigma(y)) \\ &= h(y)m(z) + \mu(y)h(z)m(\sigma(y)). \end{aligned}$$

By replacing z by $\sigma(z)$ we obtain

$$h(y\sigma(z)) = h(y)(m \circ \sigma)(z) + \mu(y)h(\sigma(z))m(\sigma(y)),\ y, z \in M. \tag{4.16}$$

By multiplying (4.16) by $\mu(z)$ and by setting $n(z) = (m \circ \sigma)(z)\mu(z)$ we obtain the μ-sine subtraction law with the companion function n

$$\mu(z)h(y\sigma(z)) = h(y)n(z) - h(z)n(y). \tag{4.17}$$

This ends the first case.
Second case : Suppose $g(e) \neq 0$. Then we obtain from (4.7) with $x = e$ that

$$h(yz) = [\mu(y)g(\sigma(y))h(z) + g(z)h(y)]/g(e). \tag{4.18}$$

Interchanging y and z in (4.18) we get

$$h(zy) = [\mu(z)g(\sigma(z))h(y) + g(y)h(z)]/g(e). \tag{4.19}$$

By replacing z by $\sigma(z)$ (4.19) and multiplying (4.19) by $\mu(z)$ we get

$$\mu(z)h(\sigma(z)y) = h(y)\frac{g}{g(e)}(z) - \frac{g}{g(e)}(y)h(z),\ y, z \in M. \tag{4.20}$$

Since h is central it follows that h satisfies the μ-sine subtraction law with the companion function $\frac{g}{g(e)}$

$$\mu(z)h(y\sigma(z)) = h(y)\frac{g}{g(e)}(z) - \frac{g}{g(e)}(y)h(z),\ y, z \in M. \tag{4.21}$$

□

In the next two theorems we obtain the solutions of equation (1.9) by using our results for the μ-sine subtraction law. We will follow the method used in [8].
Let $\mathcal{N}_\mu(\sigma, S)$ be the nullspace given by

$$\mathcal{N}_\mu(\sigma, S) = \{\theta : S \longrightarrow \mathbb{C}\ : \theta(xy) - \mu(y)\theta(\sigma(y)x) = 0,\ x, y \in G\}.$$

In the next theorem we consider the group case

Theorem 4.2. *Let G be a group, let σ be a involutive automorphism on G, let $\mu : G \longrightarrow \mathbb{C}$ be a multiplicative function such that $\mu(x\sigma(x)) = 1$ for all $x \in G$. Suppose that $f, g, h : G \longrightarrow \mathbb{C}$ satisfy functional equation (1.9). Suppose also that $g \neq 0$ and $h \neq 0$. Then there exists a character χ of G, constants c, c_1, $c_2 \in \mathbb{C}$, and a function $\theta \in \mathcal{N}_\mu(\sigma, S)$ such that one of the following holds*
i) If $\chi \neq \mu\,\chi \circ \sigma$, then

$$h = c_1\frac{\chi - \mu\,\chi \circ \sigma}{2},\ g = \frac{\chi + \mu\,\chi \circ \sigma}{2} + c_2\frac{\chi - \mu\,\chi \circ \sigma}{2},$$

$$f = \theta + \frac{c_1}{2}[c\frac{\chi - \mu\,\chi \circ \sigma}{2} + c_2\frac{\chi + \mu\,\chi \circ \sigma}{2}]$$

ii) If $\chi = \mu\,\chi\circ\sigma$, *then*

$$h = \chi A,\ \ g = \chi(c + c_2 A), f = \theta + \chi A(\frac{c}{2} + \frac{c_2}{4}A).$$

where $A : G \longrightarrow \mathbb{C}\setminus\{0\}$ *is an additive function such that* $A\circ\sigma = -A \neq 0$.
Conversely, the formulas of (i) and (ii) define solutions of (1.9).
Moreover, if G *is a topological group, and and* $f, g, h \in \mathcal{C}(G)$, *then* $\chi, \mu\,\chi\circ\sigma, A, \theta \in \mathcal{C}(G)$, *while* $A \in \mathcal{C}(G)$.

The proof of the theorem 4.2 will be integrated into that of theorem 4.3 in which we consider the monoid case

Theorem 4.3. *Let* M *be a monoid which is generated by its squares, let* σ *be an involutive automorphism on* M, *let* $\mu : M \longrightarrow \mathbb{C}$ *be a multiplicative function such that* $\mu(x\sigma(x)) = 1$ *for all* $x \in M$. *Suppose that* $f, g, h : M \longrightarrow \mathbb{C}$ *satisfy functional equation (1.9). Suppose also that* $g \neq 0$ *and* $h \neq 0$. *Then there exists a multiplicative function* $\chi : M \longrightarrow \mathbb{C}$ $\chi \neq 0$, *constants* c, c_1, $c_2 \in \mathbb{C}$, *and a function* $\theta \in \mathcal{N}_\mu(\sigma, S)$ *such that one of the following holds*
i) If $\chi \neq \mu\,\chi\circ\sigma$, *then*

$$h = c_1\frac{\chi - \mu\,\chi\circ\sigma}{2},\ g = \frac{\chi + \mu\,\chi\circ\sigma}{2} + c_2\frac{\chi - \mu\,\chi\circ\sigma}{2},$$

$$f = \theta + \frac{c_1}{2}[c\frac{\chi - \mu\,\chi\circ\sigma}{2} + c_2\frac{\chi + \mu\,\chi\circ\sigma}{2}]$$

ii) If $\chi = \mu\,\chi\circ\sigma$, *then* $h(x) = g(x) = 0$ *and* $f(x) = \theta(x)$ *for* $x \in I_\chi$, *and*

$$h(x) = \chi(x)A(x),\ \ g(x) = \chi(x)(c + c_2 A(x)), f(x) = \theta(x) + \chi(x)A(x)(\frac{c}{2} + \frac{c_2}{4}A(x))$$

for $x \in M\setminus I_\chi$ *where* $A : M\setminus I_\chi \longrightarrow \mathbb{C}\setminus\{0\}$ *is an additive function such that* $A\circ\sigma = -A \neq 0$.
Furthermore, if M *is a topological monoid and* $k \in C(M)$, *then* l, χ, $\mu\chi\circ\sigma$, $A \in \mathcal{C}(M)$. *Conversely, the formulas of (i) and (ii) define solutions of (1.9).*
Moreover, if M *is a topological monoid, and* $f, g, h \in \mathcal{C}(M)$, *then* $\chi, \mu\,\chi\circ\sigma, A, \theta \in \mathcal{C}(M)$, *while* $A \in \mathcal{C}(M\setminus I_\chi)$.

Proof. According to Proposition 4.1 we have two following cases:
First case : Suppose that $g(e) = 0$, then h satisfies the μ-sine subtraction law and $g = bh$ where $b \in \mathbb{C}\setminus\{0\}$. According to Theorem 2.1 and Proposition 2.2 we get (for M is a group or a monoid) that if $\chi \neq \mu\chi\circ\sigma$, then $h = c_1\frac{\chi - \mu\,\chi\circ\sigma}{2}$ where $c_1 \in \mathbb{C}$. Since $g = bh$, then $g = c_2\frac{\chi - \mu\,\chi\circ\sigma}{2}$ where $c_2 \in \mathbb{C}$. Subsisting g and h in (3.1) we get for all $x, y \in M$

$$\begin{aligned} f(xy) - \mu(y)f(\sigma(y)x) &= g(x)h(y) \\ &= \frac{c_1c_2}{4}[\chi(x) - \mu(x)\chi(\sigma(x))][\chi(y) - \mu(y)\chi(\sigma(y))] \\ &= \frac{c_1c_2}{4}[\chi(xy) - \mu(y)\chi(\sigma(y)x) - \mu(x)\chi(\sigma(\sigma(y)x)) \\ &\quad +\mu(xy)\chi(\sigma(xy))]. \end{aligned}$$

Let $\theta = f - \frac{c_1c_2}{4}(\chi + \mu\,\chi\circ\sigma)$ we have $\theta \in \mathcal{N}_\mu(\sigma, S)$ and $f = \theta + \frac{c_1}{2}[c\frac{\chi - \mu\,\chi\circ\sigma}{2} + c_2\frac{\chi + \mu\,\chi\circ\sigma}{2}]$ with $c = 0$.
Now, if $\chi = \mu\chi\circ\sigma$.
When M is a group then we get from Theorem 2.1 that $h = \chi A$ where A is an

additive function such that $A \circ \sigma = -A \neq 0$. Since $g = bh$, then $g = b\chi A = c_2\chi A$. Subsisting g and h in (3.1) we get

$$\begin{aligned} f(xy) - \mu(y)f(\sigma(y)x) &= g(x)h(y)c_2\chi(x)A(x)\chi(y)A(y) \\ &= \frac{c_2}{4}[\chi(y)A(xy)^2 - \chi(\sigma(y))A(\sigma(y)x)^2]. \end{aligned}$$

Let $\theta = f - c_2\frac{\chi A^2}{4}$. Then $\theta \in \mathcal{N}_\mu(\sigma, S)$.
When M is a monoid, by Proposition 2.2 and and by the same way as in [6] we have $\theta = \theta_1 \cup \theta_2$ where where $\theta_1(x) = f(x) - c_2\frac{\chi(x)A^2(x)}{4}$ on $M \setminus I_\chi$ and $\theta_2(x) = f(x)$ on I_χ.
Second case : Suppose $g(e) \neq 0$. We have h and $\frac{g}{g(e)}$ play the role of k and l respectively in Theorem 2.1 or in Proposition 2.2. If $\chi \neq \mu\ \chi\circ\sigma$, then $h = c_1\frac{\chi - \mu\ \chi\circ\sigma}{2}$ where $c_1 \in \mathbb{C}$ and $g = c\frac{\chi+\mu\ \chi\circ\sigma}{2} + c_2\frac{\chi-\mu\ \chi\circ\sigma}{2}$. By the same way as in [9] we get that $\theta = f - c_1\frac{[(c+c_2)\chi-(c-c_2)\chi\circ\sigma]}{4} \in \mathcal{N}_\mu(\sigma, S)$.
Finally, if $g(e) \neq 0$ and $\chi = \mu\ \chi \circ \sigma$ we get the remainder by the same way as in [9]. □

5. Applications: Solutions of equation (1.11) on groups and monoids

In this section, we use the results obtained in the previous paragraph to solve the functional equation (1.11) on groups and monoids. We proceed as follows to reduce the equation to the functional equation (1.8) and (1.9) so that we can apply Theorem 3.1, Theorem 3.2, Theorem 4.2. and Theorem 4.3.

Theorem 5.1. *Let G be a group, and σ an homomorphism involutive of G. Let $\mu : G \longrightarrow \mathbb{C}$ be a multiplicative function such that $\mu(x\sigma(x)) = 1$ for all $x \in G$. Suppose that the functions $f, g, h : G \longrightarrow \mathbb{C}$ satisfy the functional equation (1.11). Suppose also that $f+g \neq 0$. Then there exists a character χ of G, constants $\alpha \in \mathbb{C}^*$, c_1, $c_2 \in \mathbb{C}$, and a function $\theta \in \mathcal{N}_\mu(\sigma, S)$ such that one of the following holds*
(a) If $\chi \neq \mu\ \chi \circ \sigma$, then $f = \frac{1}{2}[(1 + \frac{c_1c_2}{2})\frac{\chi+\mu\ \chi\circ\sigma}{2} + 2c_2\frac{\chi-\mu\ \chi\circ\sigma}{2} + \theta]$; $g = \frac{1}{2}[(1 - \frac{c_1c_2}{2})\frac{\chi+\mu\ \chi\circ\sigma}{2} - \theta]$ and $h = \frac{1}{\alpha}[\frac{\chi+\mu\ \chi\circ\sigma}{2} + c_2\frac{\chi-\mu\ \chi\circ\sigma}{2}]$.
(b) If $\chi = \mu\ \chi \circ \sigma$ then $f = \frac{c_2\chi(2+A)+\theta+\chi Ac_2(1+\frac{A}{4})}{2}$; $g = \frac{c_2\chi(2+A)-\theta-\chi Ac_2(1+\frac{A}{4})}{2}$ and $h = \frac{1}{\alpha}c_2\chi(2 + A)$.

Proof. Let $f, g, h : G \longrightarrow \mathbb{C}$ satisfy the functional equation (1.11). The case $g = -f$ was treated in Theorem 3.1 and Theorem 3.2. From now, on we assume that $f+g \neq 0$. Let $h_o := \frac{h-\mu h\circ\sigma}{2}$ respectively $h_e := \frac{h+\mu h\circ\sigma}{2}$ denote the odd, respectively even, part of h with respect to μ and σ.
Setting $x = e$ in (1.11) gives us

$$f(y) + \mu(y)g(\sigma(y)) = h(e)h(y) \tag{5.1}$$

for all $y \in G$. Taking $y = e$ in (1.11) and using $\mu(e) = 1$ we find

$$f(x) + g(x) = h(e)h(x) \tag{5.2}$$

for all $x \in G$. So, by comparing (5.1) with (5.2) we get

$$g(x) = \mu(x)g(\sigma(x)),\ x \in G. \tag{5.3}$$

We note that $(f + g)(xy) + \mu(y)(f + g)(\sigma(y)x) = f(xy) + \mu(y)g(\sigma(y)x) + g(xy) + \mu(y)f(\sigma(y)x) = h(x)h(y) + g(xy) + \mu(y)f(\sigma(y)x)$. By using (5.3) we have $g(xy) =$

$\mu(xy)g(\sigma(x)\sigma(y))$, then we get $g(xy)+\mu(y)f(\sigma(y)x) = \mu(y)f(\sigma(y)x)+\mu(xy)g(\sigma(x)\sigma(y)) = \mu(y)[f(\sigma(y)x) + \mu(x)g(\sigma(x)\sigma(y))] = \mu(y)h(x)h(\sigma(y))$. Which implies that

$$(f+g)(xy) + \mu(y)(f+g)(\sigma(y)x) = 2h(x)h_e(y) \tag{5.4}$$

for all $x, y \in G$. From (5.2) and the assumption that $f+g \neq 0$ we get $h(e) \neq 0$. So, equation (5.4) can be written as follows

$$(f+g)(xy) + \mu(y)(f+g)(\sigma(y)x) = 2(f+g)(x)\frac{h_e(y)}{h(e)} \tag{5.5}$$

for all $x, y \in G$.
On the other hand by using similar computation used above, we obtain

$$(f-g)(xy) - \mu(y)(f-g)(\sigma(y)x) = 2h(x)h_o(y) = (f+g)(x)\frac{2h_o(y)}{h(e)} \tag{5.6}$$

for all $x, y \in G$. We can now apply Theorem 3.1, Theorem 3.2, Theorem 4.2 and Theorem 4.3.
If $h_o = 0$, then $f - g \in \mathcal{N}(\sigma, G)$, so there exists $\theta \in \mathcal{N}_\mu(\sigma, G)$ such that $f - g = \theta$. Since f, g satisfy (5.5) then from Theorem 3.1 we get the only possibility $f + g = \alpha^2 \frac{\chi+\mu\,\chi\circ\sigma}{2}$ and $h = \alpha\frac{\chi+\mu\,\chi\circ\sigma}{2}$ for some character $\chi : G \longrightarrow \mathbb{C}$ and a constant $\alpha \in \mathbb{C}$ and we deduce that $f = \frac{1}{2}[\theta + \alpha^2(\frac{\chi+\mu\,\chi\circ\sigma}{2})]$; $g = \frac{1}{2}[-\theta + \alpha^2(\frac{\chi+\mu\,\chi\circ\sigma}{2})]$. We deal with case (i).
So during the rest of the proof we will assume that $h_o \neq 0$. the function $f + g$, h_o are solution of equation (5.6) with $f+g \neq 0$ and $h_o \neq 0$, so we know from Theorem 4.2 that there are only the following 2 possibilities:
(i) $f - g = \theta + \frac{c_1}{2}[c\frac{\chi-\mu\,\chi\circ\sigma}{2} + c_2\frac{\chi+\mu\,\chi\circ\sigma}{2}]$; $f + g = \frac{\chi+\mu\,\chi\circ\sigma}{2} + c_2\frac{\chi-\mu\,\chi\circ\sigma}{2}$ for some character χ on G such that $\chi \neq \mu\,\chi\circ\sigma$, $\theta \in \mathcal{N}_\mu(\sigma, G)$ and constants $c, c_1, c_2 \in \mathbb{C}$. So, we have $g = \frac{1}{2}[(1 - \frac{c_1c_2}{2})\frac{\chi+\mu\,\chi\circ\sigma}{2} + (c_2 - \frac{cc_1}{2})\frac{\chi-\mu\,\chi\circ\sigma}{2} - \theta]$; $f = \frac{1}{2}[(1 + \frac{c_1c_2}{2})\frac{\chi+\mu\,\chi\circ\sigma}{2} + (c_2 + \frac{cc_1}{2})\frac{\chi-\mu\,\chi\circ\sigma}{2} + \theta]$. Since $g = \mu g \circ \sigma$, then we have $c_2 = \frac{cc_1}{2}$ and f, g are as follows: $f = \frac{1}{2}[(1 + \frac{c_1c_2}{2})\frac{\chi+\mu\,\chi\circ\sigma}{2} + 2c_2\frac{\chi-\mu\,\chi\circ\sigma}{2} + \theta]$; $g = \frac{1}{2}[(1 - \frac{c_1c_2}{2})\frac{\chi+\mu\,\chi\circ\sigma}{2} - \theta]$. We deal with case (a).
(ii) $f - g = \theta + \chi A(\frac{c}{2} + \frac{c_2}{4}A)$; $f + g = \chi(c + c_2A)$, where χ is a character on G such that $\chi = \mu\,\chi\circ\sigma$, A is an additive map on G such that $A\circ\sigma = -A$, $\theta \in \mathcal{N}_\mu(\sigma, G)$ and $c, c_2 \in \mathbb{C}$. So, we get $g = \frac{\chi(c+c_2A)-\theta-\chi A(\frac{c}{2}+\frac{c}{4})A}{2}$. Since $g = \mu g \circ \sigma$ so, we have $c = 2c_2$. Consequently we have $g = \frac{c_2\chi(2+A)-\theta-\chi Ac_2(1+\frac{A}{4})}{2}$; $f = \frac{c_2\chi(2+A)+\theta+\chi Ac_2(1+\frac{A}{4})}{2}$, where $A : G \longrightarrow \mathbb{C}$ is an additive function such that $A \circ \sigma(x) = -A(x)$ for all $x \in G$. We deal with case (b). □

Theorem 5.2. *Let M be a monoid which is generated by its squares, let σ an involutive automorphism on M. Let $\mu : M \longrightarrow \mathbb{C}$ be a multiplicative function such that $\mu(x\sigma(x)) = 1$ for all $x \in M$. Suppose that $f, g, h : M \longrightarrow \mathbb{C}$ satisfy the functional equation (1.11). Suppose also that $f + g \neq 0$. Then there exists a character χ of M, constants $\alpha \in \mathbb{C}^*$, c_1, $c_2 \in \mathbb{C}$, and a function $\theta \in \mathcal{N}_\mu(\sigma, M)$ such that one of the following holds*
(a) If $\chi \neq \mu\,\chi\circ\sigma$, then $f = \frac{1}{2}[(1 + \frac{c_1c_2}{2})\frac{\chi+\mu\,\chi\circ\sigma}{2} + 2c_2\frac{\chi-\mu\,\chi\circ\sigma}{2} + \theta]$; $g = \frac{1}{2}[(1 - \frac{c_1c_2}{2})\frac{\chi+\mu\,\chi\circ\sigma}{2} - \theta]$ and $h = \frac{1}{\alpha}[\frac{\chi+\mu\,\chi\circ\sigma}{2} + c_2\frac{\chi-\mu\,\chi\circ\sigma}{2}]$.
(b) If $\chi = \mu\,\chi\circ\sigma$ then $f(x) = \frac{\theta(x)}{2}$, $g(x) = \frac{-\theta(x)}{2}$ and $h(x) = 0$ for all $x \in I_\chi$; $f(x) = \frac{c_2\chi(2+A(x))+\theta(x)+\chi(x)Ac_2(1+\frac{A(x)}{4})}{2}$; $g(x) = \frac{c_2\chi(2+A(x))-\theta(x)-\chi(x)Ac_2(1+\frac{A(x)}{4})}{2}$

and $h(x) = \frac{1}{\alpha}c_2\chi(2 + A(x))$ *for all* $x \in M\backslash I_\chi$ *and where* $A : M\backslash I_\chi \longrightarrow \mathbb{C}$ *is an additive function such that* $A \circ \sigma(x) = -A(x)$ *for all* $x \in M\backslash I_\chi$.

References

[1] Aczél, J., Dhombres, J. Functional equations in several variables. With applications to mathematics, information theory and to the natural and social sciences. Encyclopedia of Mathematics and its Applications, vol. 31. Cambridge University Press, Cambridge, 1989

[2] d'Alembert, J. Recherches sur la courbe que forme une corde tendue mise en vibration, I. Hist. Acad. Berlin 1747(1747), 214-219

[3] d'Alembert, J.: Recherches sur la courbe que forme une corde tendue mise en vibration, II. Hist. Acad. Berlin 1747(1747), 220-249

[4] d'Alembert, J. Addition au Mémoire sur la courbe que forme une corde tendue mise en vibration. Hist. Acad. Berlin 1750(1750), 355-360

[5] Davison, Thomas M.K. D'Alembert's functional equation on topological monoids, Publ. Math. Debrecen 75(2009), 41-66

[6] Ebanks, B., Stetkær, H. On Wilson's functional equations, Aequationes Math., 89(2015), 339-354

[7] Ebanks, B.R., Sahoo, P.K., Sander, W. Characterizations of Information Measures. World Scientific Publishing Co, Singapore, 1998

[8] Ebanks, B.R., Stetkær, H. d'Alembert's other functional equation on monoids with involution. Aequationes Math. 89(2015), 187-206

[9] Elqorachi, E., Redouani, A. Solutions and stability of a variant of Wilson's functional equation, arXiv:1505.06512v1 [math.CA], 2015

[10] Parnami, J.C, Singh, H., Vasudeva, H.L. On an exponential-cosine functional equation. Period. Math. Hungar. 19(1988), 287-297

[11] Stetkær, H.: On a variant of Wilson's functional equation on groups, Aequationes Math., 68(2004), 160-176

[12] Stetkær, H. Functional equations on abelian groups with involution. Aequationes Math. 54(1997), 144-172

[13] Stetkær, H. Functional Equations on Groups. World Scientific Publishing Co, Singapore, 2013

[14] Stetkær, H. A variant of d'Alembert's functional equation, Aequationes Math., 89(2015), 657-662

[15] Stetkær, H. d'Alembert's functional equation on groups. Recent developments in functional equations and inequalities, pp. 173-191, Banach Center Publ., vol. 99. Polish Acad. Sci. Inst. Math., Warsaw, 2013

[16] Yang, D. The symmetrized Sine addition formula, Aequationes Math., 82(2011), 299-318

Belaid Bouikhalene
Departement of Mathematics and Informatics
Polydisciplinary Faculty, Sultan Moulay Slimane university, Beni Mellal, Morocco.
E-mail : bbouikhalene@yahoo.fr.

Elhoucien Elqorachi,
Department of Mathematics,
Faculty of Sciences, Ibn Zohr University, Agadir, Morocco,
E-mail: elqorachi@hotmail.com